

Clinical Policy: Adagrasib (Krazati)

Reference Number: CP.PHAR.605

Effective Date: 03.01.23 Last Review Date: 02.25

Line of Business: Commercial, HIM, Medicaid Revision Log

See <u>Important Reminder</u> at the end of this policy for important regulatory and legal information.

Description

Adagrasib (Krazati®) is an inhibitor of the RAS GTPase family.

FDA Approved Indication(s)

Krazati is indicated:

- As a single agent, for the treatment of adult patients with *KRAS* G12C-mutated locally advanced or metastatic non-small cell lung cancer (NSCLC), as determined by an FDA-approved test, who have received at least one prior systemic therapy*.
- In combination with cetuximab, for the treatment of adult patients with *KRAS* G12C-mutated locally advanced or metastatic colorectal cancer (CRC), as determined by an FDA-approved test, who have received prior treatment with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy*.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation® that Krazati is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

- A. Non-Small Cell Lung Cancer (must meet all):
 - 1. Diagnosis of recurrent, advanced, or metastatic NSCLC;
 - 2. Prescribed by or in consultation with an oncologist;
 - 3. Age \geq 18 years;
 - 4. For Krazati requests, member must use adagrasib, if available, unless contraindicated or clinically significant adverse effects are experienced;
 - 5. Disease is positive for *KRAS* G12C mutation;
 - 6. Member meets one of the following (a or b):
 - a. Both of the following (i and ii):
 - i. Member has received at least one systemic therapy (not including Lumakras) (see Appendix B);
 - ii. Member must use Lumakras®, unless contraindicated, clinically significant adverse effects are experienced, or request is for treatment associated with

^{*}This indication is approved under accelerated approval based on objective response rate (ORR) and duration of response (DOR). Continued approval for this indication may be contingent upon verification and description of a clinical benefit in a confirmatory trial(s).



cancer for a State with regulations against step therapy in certain oncology settings (see Appendix D);

- b. Member has brain metastases;
- 7. Prescribed as monotherapy;
- 8. Request meets one of the following (a or b):*
 - a. Dose does not exceed 1,200 mg (6 tablets) per day;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*Prescribed regimen must be FDA-approved or recommended by NCCN

Approval duration:

Medicaid/HIM – 6 months

Commercial – 12 months or duration of request, whichever is less

B. Colorectal Cancer (must meet all):

- 1. Diagnosis of locally advanced or metastatic CRC;
- 2. Prescribed by or in consultation with an oncologist;
- 3. Age \geq 18 years;
- 4. For Krazati requests, member must use adagrasib, if available, unless contraindicated or clinically significant adverse effects are experienced;
- 5. Disease is positive for *KRAS* G12C mutation;
- 6. Member has had previous treatment with a fluoropyrimidine- (e.g., 5-fluorouracil, capecitabine), oxaliplatin-, and irinotecan-containing chemotherapy (*see Appendix B*);
- 7. Krazati is prescribed in one of the following ways (a, b, or c):
 - a. In combination with Vectibix® (off-label);
 - b. In combination with Erbitux[®];
 - c. As monotherapy if member is unable to tolerate epidermal growth factor receptor (EGFR) inhibitor (e.g., Erbitux, Vectibix) due to toxicity;
- 8. Request meets one of the following (a or b):*
 - a. Dose does not exceed 1,200 mg (6 tablets) per day;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*Prescribed regimen must be FDA-approved or recommended by NCCN

Approval duration:

Medicaid/HIM – 6 months

Commercial – 12 months or duration of request, whichever is less

C. NCCN Compendium Indications (off-label) (must meet all):

- 1. Diagnosis of one of the following (a, b, or c):
 - a. Pancreatic adenocarcinoma that is locally advanced, recurrent, or metastatic;
 - b. Ampullary adenocarcinoma;
 - c. One of the following biliary tract cancers that is unresectable, resected gross residual (R2), or metastatic (i, ii, or iii) as subsequent therapy:
 - i. Extrahepatic cholangiocarcinoma;
 - ii. Intrahepatic cholangiocarcinoma;
 - iii. Gallbladder cancer;
- 2. Prescribed by or in consultation with an oncologist;
- 3. Age \geq 18 years;



- 4. For Krazati requests, member must use adagrasib, if available, unless contraindicated or clinically significant adverse effects are experienced;
- 5. Prescribed as monotherapy;
- 6. Disease is positive for *KRAS* G12C mutation;
- 7. Request meets one of the following (a or b):*
 - a. Dose does not exceed 1,200 mg (6 tablets) per day;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*Prescribed regimen must be FDA-approved or recommended by NCCN

Approval duration:

Medicaid/HIM – 6 months

Commercial – 12 months or duration of request, whichever is less

D. Other diagnoses/indications (must meet 1 or 2):

- 1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business:
 CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
- 2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. All Indications in Section I (must meet all):

- 1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving Krazati for a covered indication and has received this medication for at least 30 days;
- 2. Member is responding positively to therapy;
- 3. For Krazati requests, member must use adagrasib, if available, unless contraindicated or clinically significant adverse effects are experienced;
- 4. If request is for a dose increase, request meets one of the following (a or b):*
 - a. New dose does not exceed 1,200 mg (6 tablets) per day;
 - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*Prescribed regimen must be FDA-approved or recommended by NCCN

Approval duration:

Medicaid/HIM – 12 months

Commercial – 12 months or duration of request, whichever is less



B. Other diagnoses/indications (must meet 1 or 2):

- 1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business:
 CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
- 2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

CRC: colorectal cancer FDA: Food and Drug Administration EGFR: epidermal growth factor receptor NSCLC: non-small cell lung cancer

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business

and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Lumakras (sotorasib)	NSCLC:	960 mg/day
	960 mg PO QD	
cisplatin- or carboplatin-	NSCLC:	Varies
containing chemotherapy	Varies	
Imfinzi [®]	NSCLC:	1,500 mg every 4
(durvalumab)	10 mg/kg IV every 2 weeks or 1,500 mg	weeks
	every 4 weeks	
Keytruda®	NSCLC:	400 mg every 6
(pembrolizumab)	200 mg IV every 3 weeks OR 400 mg	weeks
	every 6 weeks up to 24 months	



Drug Name	Dosing Regimen	Dose Limit/
		Maximum Dose
Libtayo®	NSCLC:	350 mg every 3
(cemiplimab-rwlc)	350 mg IV every 3 weeks	weeks
Opdivo®	NSCLC:	480 mg every 4
(nivolumab)	240 mg IV every 2 weeks or 480 mg IV	weeks
	every 4 weeks	
Tecentriq®	NSCLC:	1,680 mg every 4
(atezolizumab)	840 mg IV every 2 weeks, 1,200 mg IV	weeks
	every 3 weeks, or 1,680 mg IV every 4	
	weeks	
Yervoy®	NSCLC:	1 mg/kg every 6
(ipilimumab)	In combination with Opdivo: 1 mg/kg IV	weeks
	every 6 weeks	
Imjudo® (tremelimumab)	NSCLC:	See dosing
	Patients \geq 30 kg: 75 mg IV on day 1	regimen
	every 3 weeks for cycles 1 through 4 in	
	combination with durvalumab and	
	platinum-based chemotherapy; cycle 5 no	
	dose; cycle 6 - 75 mg IV on day 1 in	
	combination with durvalumab	
	Patients < 30 kg: 1 mg/kg on day 1 every	
	3 weeks for cycles 1 through 4 in	
	combination with durvalumab and	
	platinum-based chemotherapy; cycle 5 no	
	dose; cycle 6 - 1 mg/kg IV on day 1 in	
	combination with durvalumab	
fluoropyrimidine-	CRC:	Varies
containing regimens (e.g.,	Varies	
5-fluorouracil,		
capecitabine, FOLFOX,		
CAPEOX, FOLFIRINOX		
oxaliplatin-containing	CRC:	Varies
regimens (e.g., FOLFOX,	Varies	
CAPEOX,		
FOLFIRINOX)		
irinotecan-containing	CRC:	Varies
regimens (e.g., FOLFIRI,	Varies	
FOLFIRINOX)		

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings None reported



Appendix D: States with Regulations against Redirections in Cancer

State	Step Therapy Prohibited?	Notes	
FL	Yes	For stage 4 metastatic cancer and associated conditions	
GA	Yes	For stage 4 metastatic cancer. Redirection does not refer to	
		review of medical necessity or clinical appropriateness	
IA	Yes	For standard of care stage 4 cancer drug use, supported by peer-	
		reviewed, evidence-based literature, and approved by FDA	
LA	Yes	For stage 4 advanced, metastatic cancer or associated conditions.	
		Exception if "clinically equivalent therapy, contains identical	
		active ingredient(s), and proven to have same efficacy	
MS	Yes	*Applies to HIM requests only*	
		For advanced metastatic cancer and associated conditions	
NV	Yes	Stage 3 and stage 4 cancer patients for a prescription drug to treat	
		the cancer or any symptom thereof of the covered person	
OH	Yes	*Applies to Commercial and HIM requests only*	
		For stage 4 metastatic cancer and associated conditions	
OK	Yes	*Applies to HIM requests only*	
		For advanced metastatic cancer and associated conditions	
PA	Yes	For stage 4 advanced, metastatic cancer	
TN	Yes	For advanced metastatic cancer and associated conditions	
TX	Yes	For stage 4 advanced, metastatic cancer and associated conditions	

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
NSCLC, CRC	600 mg PO BID	1,200 mg/day

VI. Product Availability

Tablet: 200 mg

VII. References

- 1. Krazati Prescribing Information. San Diego, CA: Mirati Therapeutics, Inc.; July 2024. Available at: https://packageinserts.bms.com/pi/pi krazati.pdf. Accessed October 31, 2024.
- 2. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at http://www.nccn.org. Accessed November 11, 2024.
- 3. National Comprehensive Cancer Network. Non-Small Cell Lung Cancer Version 11.2024. Available at: http://www.nccn.org/professionals/physician_gls/pdf/nscl.pdf. Accessed November 11, 2024.
- 4. National Comprehensive Cancer Network. Colon Cancer Version 5.2024. Available at: http://www.nccn.org/professionals/physician_gls/pdf/colon.pdf. Accessed November 11, 2024.
- 5. National Comprehensive Cancer Network. Rectal Cancer Version 4.2024. Available at: http://www.nccn.org/professionals/physician_gls/pdf/rectal.pdf. Accessed November 11, 2024.
- 6. Jänne PA, Riely GJ, Gadgeel SM, et al. Adagrasib in non-small-cell lung cancer harboring a *KRASG12C* mutation. N Engl J Med. 2022 Jul 14;387(2):120-131.



Reviews, Revisions, and Approvals	Date	P&T
		Approval
		Date
Policy created.	01.04.23	02.23
1Q 2024 annual review: Per NCCN: for NSCLC added	10.26.23	02.24
monotherapy criterion and added brain metastases as exception to		
having received ≥ 1 prior therapy; added off-label criteria for		
pancreatic adenocarcinoma and colorectal cancer; references		
reviewed and updated.		
RT4: added new FDA-approved indication for the treatment of	06.27.24	
CRC; updated Appendix B.		
Ad hoc: added off-label criteria for ampullary adenocarcinoma and		
biliary tract cancers as subsequent therapy per NCCN compendium.		
1Q 2025 annual review: added generic redirection for initial and	12.02.24	02.25
continued therapy per template; per NCCN, for NSCLC removed		
"locally" from "locally advanced" NSCLC option and for CRC		
added monotherapy option; references reviewed and updated. Per		
December SDC: for NSCLC, added redirection to Lumakras for		
members without brain metastases; for all indications, removed		
requirement that member has not received prior treatment with		
Lumakras.		

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to



applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2023 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.