

Preemptive policy: This is a P&T approved policy and can be used after the drug is FDA approved until it is superseded by an updated policy



Clinical Policy: Tiratricol (Emcitate)

Reference Number: CP.PHAR.790

Effective Date: **FDA Approval Date**

Last Review Date: 08.26

Line of Business: Commercial, HIM/ICHRA, Medicaid

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Tiratricol (Emcitate^{®/TM}) is a thyroid hormone analogue.

FDA Approved Indication(s) **[Pending]**

Emcitate is indicated for the treatment of patients with monocarboxylate transporter 8 (MCT8) deficiency.

Limitation(s) of use: **[XXX]**

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Emcitate is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria*

**Criteria will mirror the clinical information from the prescribing information once FDA-approved*

A. Monocarboxylate Transporter 8 Deficiency (must meet all):

1. Diagnosis of MCT8 deficiency confirmed by both of the following (a and b):*
 - a. Genetic testing confirming the presence of *SLC16A2* gene mutation;
 - b. Clinical symptoms consistent with MCT8 deficiency (e.g., neurodevelopmental delays, motor disability, poor weight gain; *see Appendix D*);
2. Prescribed by or in consultation with an endocrinologist, neurologist, geneticist, or metabolic specialist;*
3. Documentation of serum total T3 levels at baseline;*
4. Dose does not exceed **FDA maximum dose**.*

Approval duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace/ICHRA) or PDL (Medicaid), the no coverage criteria policy for the relevant line of

- business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace/ICHRA, and CP.PMN.255 for Medicaid; or
- b. For drugs NOT on the formulary (commercial, health insurance marketplace/ICHRA) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace/ICHRA, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace/ICHRA, and CP.PMN.53 for Medicaid.

II. Continued Therapy*

**Criteria will mirror the clinical information from the prescribing information once FDA-approved*

A. Monocarboxylate Transporter 8 Deficiency (must meet all):

1. Member meets one of the following (a or b):
 - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
 - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (*refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B*);
2. Member is responding positively to therapy as evidenced by decrease in serum total T3 levels from baseline;*
3. If request is for a dose increase, new dose does not exceed **FDA maximum dose**.*

Approval duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace/ICHRA) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace/ICHRA, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace/ICHRA) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace/ICHRA, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace/ICHRA, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace/ICHRA, and CP.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

FDA: Food and Drug Administration

MCT8: monocarboxylate transporter 8

Appendix B: Therapeutic Alternatives

Not applicable

Appendix C: Contraindications/Boxed Warnings [Pending]

- Contraindication(s): **pending**
- Boxed warning(s): **pending**

Appendix D: General Information

- Clinical symptoms of MCT8 deficiency
 - Motor disability
 - Severe intellectual disability
 - Severe developmental delay
 - Dystonia
 - Hypotonia
 - Low body weight
 - Hypermetabolic state
 - Tachycardia
 - Muscle wasting/ muscle weakness
 - Kinesigenic dyskinesias

V. Dosage and Administration [Pending]

Indication	Dosing Regimen	Maximum Dose
MCT8*	Pending	Pending

VI. Product Availability [Pending]

Tablet: 350 mg*

VII. References

1. ClinicalTrials.gov. Withdrawal of tiratricol treatment in males with monocarboxylate transporter 8 deficiency (MCT8 Deficiency) (ReTRIACt). Available at: <https://clinicaltrials.gov/study/NCT05579327>. Accessed May 26, 2026.
2. ClinicalTrials.gov. Triac trial II in MCT8 deficiency patients. Available at: <https://clinicaltrials.gov/study/NCT02396459>. Accessed May 26, 2026.
3. ClinicalTrials.gov. Thyroid hormone analog therapy in CMT8 deficiency: Triac trial patients. Available at: <https://clinicaltrials.gov/study/NCT02060474>. Accessed May 26, 2026.

4. Groeneweg S, Peeters RP, Moran C, et al. Effectiveness and safety of the tri-iodothyronine analogue Triac in children and adults with MCT8 deficiency: An international, single-arm, open-label, phase 2 trial. *Lancet Diabetes Endocrinol.* 2019;7(9):695–706. doi:10.1016/S2213-8587(19)30155-X.
5. Van Geest FS, Groeneweg S, van den Akker ELT, et al. Long-term efficacy of T3 analogue Triac in children and adults with MCT8 deficiency: A real-life retrospective cohort study. *J Clin Endocrinol Metab.* 2022;107(3):e1136–e1147. doi:10.1210/clinem/dgab750.
6. Persani L, Rodien P, Moran C, et al. 2024 European Thyroid Association guidelines on diagnosis and management of genetic disorders of thyroid hormone transport, metabolism and action. *Eur Thyroid J.* 2024;13(4):e240125. doi:10.1530/ETJ-24-0125.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created pre-emptively	06.16.26	08.26

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2026 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.